

10th International workshop on quantum phase transitions in nuclei and many-body systems
Dubrovnik, Croatia, July 2022

Coexistence of nuclear shapes: mean-field and beyond

Tamara Nikšić
University of Zagreb




Universal theory framework: Energy Density Functionals



- ✓ the nuclear many-body problem is effectively mapped onto a one-body problem without explicitly involving internucleon interactions





- ✓ the exact density functional is approximated with powers and gradients of ground state densities and currents



- ✓ universal density functionals can be extended from relatively light systems to superheavy nuclei and from the valley of stability to the particle drip line



- ✓ the coupling parameters of the EDF are fine-tuned to empirical data





- ✓ covariant EDFs – built from densities and currents bilinear in the Dirac spinor field of the nucleon




Basic implementation: self-consistent mean-field method

- produces energy surfaces as functions of intrinsic deformation parameters



- includes static correlations: deformations and pairing
- does not include collective correlations originating from symmetry restoration and quantum fluctuations around mean-field minima

octupole deformation




Beyond mean-field correlations: GCM

Restoration of broken symmetries
(rotational, parity, particle number)
and fluctuations of collective variables
(quadrupole deformation)

1. Constraint mean-field calculation


2. Angular momentum, parity and
particle number projection

3. Configuration mixing (generator
coordinate method)



Beyond mean-field correlations: Collective Hamiltonian

... nuclear excitations determined by quadrupole vibrational and rotational degrees of freedom




$$H_{\text{coll}} = \mathcal{T}_{\text{vib}}(\beta, \gamma) + \mathcal{T}_{\text{rot}}(\beta, \gamma, \Omega) + \mathcal{V}_{\text{coll}}(\beta, \gamma)$$

$$\mathcal{T}_{\text{vib}} = \frac{1}{2} B_{\beta\beta} \dot{\beta}^2 + \beta B_{\beta\gamma} \dot{\beta} \dot{\gamma} + \frac{1}{2} \beta^2 B_{\gamma\gamma} \dot{\gamma}^2$$

$$\mathcal{T}_{\text{rot}} = \frac{1}{2} \sum_{k=1}^3 \mathcal{I}_k \omega_k^2$$

The entire dynamics of the collective Hamiltonian is governed by the seven functions of the intrinsic deformations β and γ : the collective potential, the three mass parameters: $B_{\beta\beta}$, $B_{\beta\gamma}$, $B_{\gamma\gamma}$, and the three moments of inertia \mathcal{I}_k .


... collective eigenfunction:
$$\Psi_{\alpha}^{IM}(\beta, \gamma, \Omega) = \sum_{K \in \Delta I} \psi_{\alpha K}^I(\beta, \gamma) \Phi_{MK}^I(\Omega)$$



✓ an intuitive interpretation of mean-field results in terms of *intrinsic shapes* and *single-particle states*

✓ the *full model space* of occupied states can be used; no distinction between core and valence nucleons, *no need for effective charges!*


Phys. Rev. C 89, 054306 (2014)



Global analysis of quadrupole shape invariants

Phys. Rev. C 95, 054321 (2017)

- Systematic analysis of characteristic signatures of coexisting shapes in different mass regions
- Calculation includes 621 even-even nuclei with $Z, N > 10$ and for which 2_1^+ state has been determined in experiment



Global analysis of quadrupole shape invariants

Phys. Rev. C 95, 054321 (2017)

The lowest-order quadrupole invariants:

$$q_2(0_i^+) = \sum_j \langle 0_i^+ || Q || 2_j^+ \rangle \langle 2_j^+ || Q || 0_i^+ \rangle.$$

$$q_3(0_i^+) = \sqrt{\frac{7}{10}} \sum_{jk} \langle 0_i^+ || Q || 2_j^+ \rangle \langle 2_j^+ || Q || 2_k^+ \rangle \langle 2_k^+ || Q || 0_i^+ \rangle.$$

Deformation parameters:

$$q_2(0_i^+) = \left(\frac{3ZeR^2}{4\pi} \right)^2 \langle \beta^2 \rangle \equiv \left(\frac{3ZeR^2}{4\pi} \right)^2 \beta_{eff}^2 \quad R = r_0 A^{1/3}$$

$$\frac{q_3(0_i^+)}{q_2^{3/2}(0_i^+)} = \frac{\langle \beta^3 \cos 3\gamma \rangle}{\langle \beta^2 \rangle^{3/2}} \equiv \cos 3\gamma_{eff}$$



$$r_0 = 1.2 \text{ fm}$$

Global analysis of quadrupole shape invariants

Phys. Rev. C 95, 054321 (2017)

Signatures of shape coexistence:



- The difference between $\beta_{\text{eff}} \cos(3\gamma_{\text{eff}})$ for the two lowest 0^+ states is large
- The excitation energy of 0_2^+ is low in comparison to the excitation energy of the 2_1^+ state



Global analysis of quadrupole shape invariants





Phys. Rev. C 95, 054321 (2017)

- $Z \approx 64$ and $N \approx 76$: medium deformed triaxial ground state coexisting with highly deformed prolate excited state






Coexisting shapes in neutron-deficient Nd and Sm isotopes

Phys. Rev. C 98, 054308 (2018)




Coexisting shapes in neutron-deficient Nd and Sm isotopes

Phys. Rev. C 98, 054308 (2018)




Coexisting shapes in neutron-deficient Nd and Sm isotopes

Phys. Rev. C 98, 054308 (2018)




Quadrupole and octupole shape transition in thorium






...quadrupole-octupole collective Hamiltonian:

$$H_{coll} = -\frac{\hbar^2}{2\sqrt{w\mathcal{I}}} \left[\frac{\partial}{\partial\beta_2} \sqrt{\frac{\mathcal{I}}{w}} B_{33} \frac{\partial}{\partial\beta_2} - \frac{\partial}{\partial\beta_2} \sqrt{\frac{\mathcal{I}}{w}} B_{23} \frac{\partial}{\partial\beta_3} - \frac{\partial}{\partial\beta_3} \sqrt{\frac{\mathcal{I}}{w}} B_{23} \frac{\partial}{\partial\beta_2} + \frac{\partial}{\partial\beta_3} \sqrt{\frac{\mathcal{I}}{w}} B_{22} \frac{\partial}{\partial\beta_3} \right] + \frac{\hat{j}^2}{2\mathcal{I}} + V(\beta_2, \beta_3)$$

... systematics of energy spectra of the positive-parity ground-state band ($K^\pi = 0^+$) and the lowest negative-parity ($K^\pi = 0^-$) sequences in $^{224-232}\text{Th}$.





Summary

- ✓ NEDFs provide an economic, global and accurate microscopic approach to nuclear structure that can be extended from relatively light systems to superheavy nuclei, and from the valley of β -stability to the particle drip-lines.
- ✓ NEDF-based structure models that take into account collective correlations → microscopic description of low-energy observables: excitation spectra, transition rates, changes in masses, isotope and isomer shifts, related to shell evolution with nuclear deformation, angular momentum, and number of nucleons.
- ✓ NEDF-based models are applicable to large-scale calculations

Funding

This work was supported by the QuantiXLie Centre of Excellence, a project co-financed by the Croatian Government and European Union through the European Regional Development Fund - the Competitiveness and Cohesion Operational Program (Grant KK.01.1.1.01.0004).

For more information please visit:
<http://bela.phy.hr/quantixlie/hr/>
<https://strukturnifondovi.hr/>

The sole responsibility for the content of this presentation lies with the Faculty of Science, University of Zagreb. It does not necessarily reflect the opinion of the European Union.



Europska unija
Zajedno do fondova EU



Operativni program
**KONKURENTNOST
I KOHEZIJA**